
Asihma Treatment Plan — Student (This asthma action plan meets NJ Law N.J.S.A. 18A:40-12.8) (Physician's Orders)







(Please Pr	ires j			жим.рас	211j.01g			
Name			1	Date of Birth	Effective Date			
Doctor			Parent/Guardian (if applicable)		Emergency Contact	mergency Contact		
Phone			Phone		Phone	· · · · · · · · · · · · · · · · · · ·		
	(Green Zone) IIII	mor	e daily control me e effective with a	"spacer" - use i	if directed.	Triggers Check all Items that trigger		
	 Breathing is good No cough or wheeze Sleep through the night Can work, exercise, and play 	☐ Aeros ☐ Alveso ☐ Dulero ☐ Flovero ☐ Qvaro ☐ Symb ☐ Advai ☐ Asma ☐ Flovero ☐ Pulmic	r® HFA	2 puffs tw 2 puffs tw 1, 3 2 puffs tw 2 puffs tw 2 puffs tw 1, 2 500	2 puffs twice a day 2 puffs twice a day wice a day wice a day 2 puffs-twice a day 2 puffs twice a day ion twice a day ion twice a day ion twice a.day I inhalations ☐ once or ☐ twice a day bulízed ☐ once or ☐ twice a day	patient's asinma: Colds/filu Exercise Allergens Dust Mites, dust, stuffed animals, carpet Pollen - trees, grass, weeds Mold Pets - animal dander Pests - rodents, cockroaches		
And/or Peak	flow above	□ None	Remember i		fter taking inhaled medicine.	Cigarette smoke & second hand		
	If exercise triggers your		Control of the Contro	and the second state of the contract of the second state of the se	minutes before exercise.	o Perfumes,		
If quick-relief m 15-20 minutes of 2 times and syndoctor or go to And/or Peak fl	Mild wheeze Tight chest Coughing at night Other: edicine does not help within or has been used more than nptoms persist, call your the emergency room. ow from to	MEDIC Albute Xoper Albute Duon Xoper Comb Increa Other * If qu	tinue daily control me INE erol MDI (Pro-air® or Proven nex® erol □ 1.25, □ 2.5 mg eb® nex® (Levalbuterol) □ 0.31, □ pivent Respimat® ase the dose of, or add: uick-relief medicir ek, except before	dicine(s) and ADD of HOW MUCH to take an till® or Ventolin®) _2 puffs _2 puffs _1 unit _1 unit _1 unit _1 inhale _2 puffs _2 puff	nd HOW OFTEN to take it s every 4 hours as needed s every 4 hours as needed nebulized every 4 hours as needed lation 4 times a day ere than 2 times a call your doctor.	scented products. Scented products Smoke from: burning wood inside or outsi Weather Sudden temperature change Extreme weath hot and cold Ozone alert da		
Your asthma is getting worse fast: Quick-relief medicine did not help within 15-20 minutes Breathing is hard or fast Nose opens wide • Ribs show Trouble walking and talking Lips blue • Fingernails blue Other:			thma can be a life-threatening illness. Do not wait! DICINE HOW MUCH to take and HOW OFTEN to take it albuterol MDI (Pro-air® or Proventil® or Ventolin®) 4 puffs every 20 minutes (openex® 4 puffs every 20 minutes albuterol 1.25, 2.5 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 0.31, 0.63, 1.25 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 0.31, 0.63, 1.25 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.25 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.35 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.31, 1.32 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.32 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.32 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.32 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.32 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.32 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.32 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.32 mg 1 unit nebulized every 20 minutes (openex® (Levalbuterol) 1.31, 1.32		This asthma treatment plan is meant to assist not replace, the clinica decision-making required to meet			
Order him more il trans a di si rebico STADII prodella ma il soi fine. The Institutati que l'accident de la signa del si fine di rebico del collega de l'accident de la signa del si fine di rebico del collega del si fine del si fine del collega del si fine de	and the security of Milling, recognitions on testing, or deposition of the control of the contro	tudent is o proper me ebulized in ordance w student is	elf-administer Medication: apable and has been instructed athod of self-administering of the ahaled medications named above ith NJ Law. not approved to self-medicate. The parent and for physician fi	PHYSICIAN STAMP	Physician's Orders TURE	DATE		